


ZRT Iodine Test (urine)



Price: \$79.00

Short Description

Up to 10% of the population in the U.S. has iodine deficiency. Iodine is essential in healthy thyroid hormone production, brain function, breast tissue, fetal development and more.

Description

Iodine Deficiency Is a Global Health Problem

Approximately 2 billion of the world's population is iodine deficient. According to the most recent NHANES surveys, urinary iodine levels have dropped in the U.S. by half since the 1970s, and nearly 10% of the population has some degree of iodine deficiency. Iodine deficiency is resurfacing in developed countries like the U.S. and Western Europe as efforts are made to reduce the consumption of iodine-containing salt and foods (eggs and dairy) to lower blood pressure and blood lipids.

Iodine is essential in healthy thyroid hormone production, brain function, breast tissue, fetal development and much more.

Adequate Iodine is Essential at All Stages of Life

In the developing fetus, an adequate iodine supply from the mother is essential for thyroid hormone production, which is vital for proper neurological development. In those areas of the world with severe iodine deficiency, pregnant women are at increased risk of miscarriage, stillbirths, or giving birth to children with cretinism and severe neurological and developmental defects. Even in areas of the world with mild to moderate iodine deficiency, which includes some regions of the U.S., children born to mothers who are iodine deficient during pregnancy can have impaired intellectual development⁸ and are at increased risk for developing attention deficit and hyperactivity disorders.

In children and adults, iodine deficiency can lead to lower thyroid hormone synthesis and consequent symptoms and conditions of hypothyroidism. When iodine levels drop below a critical threshold, consequent low thyroid hormone levels signal the brain to increase TSH production. Excessive stimulation of the thyroid gland by TSH, caused by iodine deficiency, leads to overgrowth (goiter) in an effort to maintain steady-state thyroid hormone synthesis. Areas of the world with low iodine usually have a much higher incidence of goiter.

High Iodine Levels Can Also Be Problematic

High iodine levels, resulting from consumption of high iodine containing foods (e.g., seaweed) or iodine medications or supplements, can also disrupt normal thyroid function, causing goiter and both hyperthyroidism or hypothyroidism. Hyperthyroidism is more likely to occur in the elderly who have been exposed acutely to high iodine after living for years in a low iodine environment.

Chronic low iodine can lead to the development of autonomous thyroid nodules that produce excessive amounts of thyroid hormones (hyperthyroidism) when acutely exposed to high levels of iodine. Although rare, hyperthyroidism and associated thyrotoxicosis caused by excessive iodine consumption can cause arrhythmias and precipitate adverse cardiovascular events.

In some people, acute high iodine exposure can paradoxically suppress thyroid hormone synthesis (Wolff-Chaikoff effect) and lead to goiter and hypothyroidism.

How Much Iodine Is Needed for Optimal Thyroid Health?

Maintaining optimal iodine levels through consumption of iodine containing foods, salt, or supplements is essential for optimal thyroid hormone production. The World Health Organization (WHO), Centers for Disease Control (CDC), and other international organizations that study iodine nutritional status of world populations, unanimously agree and recommend, for non-pregnant adults, consumption of 150 µg of iodine daily for optimal thyroid function and have set 1,100 µg as the upper level of tolerance.

In Japan, where a much higher amount of iodine is consumed in the form of seaweed and fish, the Japanese Ministry of Health has set the upper limit at 3,000 µg.

The higher iodine levels found to be breast-protective are more consistent with the amounts of iodine consumed in countries like Japan, where breast cancer rate is only about one-fifth that of the U.S., and other countries with much lower iodine consumption.

Iodine's Role in Extrathyroidal Tissues

While iodine's role in thyroid hormone synthesis is well recognized, its other and equally important roles in maintaining health are less well appreciated, but well researched. Iodine serves many different and protective roles in the body including those of an antioxidant, antimicrobial, anti-proliferative, and anti-carcinogen.

Tissues other than the thyroid gland that utilize iodine include white blood cells, salivary and lacrimal glands, ciliary body of the eye, renal cortex, pancreas, liver, stomach, small and large intestinal mucosa, nasopharynx, choroid plexus, skin, adrenal cortex, mammary gland, placenta, uterus and ovary.

Iodine's ability to form iodolipids also underlies its capacity to serve as an antiproliferative and prevent abnormal benign growths (e.g., thyroid nodules, fibrocystic breasts, uterine fibroids) as well as cancers of these and other tissues (listed above).

Iodine Helps Prevent Infection

Iodine's effectiveness as an antimicrobial has been known for over a hundred years as it has been used for preoperative surgical skin preparation to prevent infection. As an antimicrobial, it is of relevance that iodine from the bloodstream concentrates in a broad spectrum of tissues that are repeatedly exposed to infectious organisms (e.g., skin, eyes, gastrointestinal tract, salivary gland, nasal tissue, breast ducts, uterine lining, and prostate).

Development of a Patient-Convenient Test to Measure Iodine Levels in Dried Urine

With iodine playing so many different roles in optimizing health and preventing disease in the thyroid and other tissues, it is essential that adequate iodine intake is maintained and problems associated with low or excessive iodine intake are identified and treated appropriately.

To this end, ZRT has developed a simple and convenient test to measure an individual's iodine level in urine dried on a filter strip. This method was developed to circumvent the more cumbersome collection of all urine over a 24 hour period.

Collecting and Testing Urinary Iodine

Iodine is collected twice during the day (first morning and last night void) on filter strips either by dipping the strip in urine collected in a cup, or by urinating directly on the strip. The urine-saturated filter strips are allowed to dry overnight and then sent to the laboratory for iodine and creatinine testing.

Advantages of Dried Urine for Iodine Testing

- Urine collection and shipment of the dried filter strips are simple and convenient for the patient and practitioner.
- Dual collections of urine directly on a filter strip, upon awakening and just before bed, are far more convenient and less subject to the inherent inaccuracies of a 24 hr urine collection.
- Iodine and creatinine in dried urine are exceptionally stable for weeks at room temperature allowing more flexibility in collection, shipment, testing, and storage.
- Iodine results expressed in μg iodine per liter of urine and per mg creatinine helps to normalize results when problems exist with urine that is very concentrated or dilute.
- Iodine testing allows for determination of iodine status based on CDC and WHO guidelines for thyroid sufficiency, as well as extra thyroidal sufficiency.

To get a thorough assessment of your iodine levels and its effect on thyroid function, see this test:

[Comprehensive Elements + Thyroid + Iodine](#)

Test Includes

Lab	
ZRT	Yes
Elements	
Iodine	Yes
Sample Type	
Blood Spot	No
Dried Urine	Yes